

## RESEARCH NOTE / NOTA CIENTÍFICA

### FREQUENCY OF ANTIBODIES AGAINST *TOXOPLASMA GONDII* IN WILD CARNIVORES AND MARSUPIALS IN NORTHEAST MEXICO

### FRECUENCIA DE ANTICUERPOS CONTRA *TOXOPLASMA GONDII* EN CARNIVOROS Y MARSUPIALES SILVESTRES EN EL NORESTE DE MEXICO

Emilio Rendón-Franco<sup>1\*</sup>; Arturo Caso<sup>2</sup>; Nayelly Gabriela Jiménez-Sánchez<sup>3</sup>;  
Sasha Carvajal-Villarreal<sup>2</sup> & Hector Zepeda-López<sup>4</sup>;

<sup>1</sup> Departamento de Producción Agrícola y Animal. Universidad Autónoma Metropolitana – Unidad Xochimilco, Calzada del Hueso 1100, Col. Villa Quietud, Delegación Coyoacán, D.F. México, C.P. 04960, México. emilio.rendon.franco@gmail.com

<sup>2</sup> Caesar Kleberg Wildlife Research Institute. Texas A&M University-Kingsville 700 University Blvd., MSC 218, Kingsville, Texas 78363-8202, Texas, USA.

<sup>3</sup> Academia de Biología. Centro de Estudios Científicos y Tecnológicos No. 6 "Miguel Othón de Mendizábal" Instituto Politécnico Nacional. Avenida Jardín y calle 4 s/n, Col. Del Gas, Delegación Azcapotzalco, D.F. México, C.P. 2950, México.

<sup>4</sup> ZELOR Instituto de Medicina de la Conservación. SAPI calle Romano No. 13525-C Fracc. Alcalá La Mesa Tijuana B.C. México C.P. 22106.

\* Correspondence should be to Emilio Rendón-Franco Calzada del Hueso 1100, Col. Villa Quietud, Delegación Coyoacán, C.P. 04960, D.F. México, 00 55 54 83 70 00 ext. 3658.

Suggested citation: Rendón-Franco, E, Caso, A, Jiménez-Sánchez, NG, Carvajal-Villarreal, S & Zepeda-López, H 2014. Frequency of antibodies against *Toxoplasma gondii* in wild carnivores and marsupials in northeast Mexico. *Neotropical Helminthology*, vol. 8, n°2, jul-dec, pp. 473-478.

#### Abstract

There is little previous information about the wild cycle of *Toxoplasma gondii* in wild carnivores and marsupials in Mexico. The objective of this study was to determine the presence of antibody against *T. gondii* in wild carnivores and marsupials in northeast Mexico. Frequency of *T. gondii* in captured specimens was: margay (*Leopardus wiedii*; n=3; 33%), jaguarundi (*Puma yagouaroundi*; n=2; 0%), bobcat (*Lynx rufus*; n= 1; 100%), coati (*Nasua narica*; n=7; 42%), coyote (*Canis latrans*; n=4; 25%), gray fox (*Urocyon cinereoargenteus*; n=3; 66%), hog-nosed skunk (*Conepatus leuconotus*; n=1; 0%), and opossum (*Didelphis* sp.; n=5; 20%). Therefore, natural infection with *T. gondii* in wild carnivores and marsupials in northeast Mexico was present.

Keywords: canidae - felidae - mustelidae - procionidae.

#### Resumen

Existe poca información acerca del ciclo silvestre de *Toxoplasma gondii* en carnívoros y marsupiales silvestres en México. El objetivo de este estudio fue determinar la presencia de anticuerpos contra *T. gondii* en carnívoros y marsupiales silvestres en el noreste de México. La frecuencia de *T. gondii* en los especímenes capturados fue: margay (*Leopardus wiedii*; n=3; 33%), jaguarundi (*Puma yagouaroundi*; n=2; 0%), lince (*Lynx rufus*; n= 1; 100%), coati (*Nasua narica*; n=7; 42%), coyote (*Canis latrans*; n=4; 25%), zorra gris (*Urocyon cinereoargenteus*; n=3; 66%), zorillo (*Conepatus leuconotus*; n=1; 0%), y tlacuache (*Didelphis* sp.; n=5; 20%). Se comprueba la infección natural con *T. gondii* en carnívoros y marsupiales silvestres en el noreste de México.

Palabras clave: canidae – felidae – mustelidae – procionidae.

## INTRODUCTION

*Toxoplasma gondii* (Nicolle, Manceaux and Splendore 1908) is an intracellular parasite that affects all homeothermic vertebrate. In the domestic cat, the complete cycle of the parasite occurs because the cat is the only definitive host.

Infected hosts may be in an infectious stage all their lives and this helps the worldwide distribution of the disease (Frenkel, 1990; Della, 1999; Wolfe, 2003). The wild felids are also considered definitive hosts since this parasite have been found in different species of wild cats (Dreesen, 1990; Wolfe, 2003; Cañon-Franco *et al.*, 2013). Even though these animals have been the definitive hosts, there are a few studies about their role on the wild cycle of the parasite. A recent revision reported that in the only neotropical small felid species that this parasite has been found is in ocelot (*Leopardus pardalis*; Linné, 1758), Geoffroy's Cat (*L. geoffroyi*; D'Orbigny and Gervais, 1844) and Oncilla (*L. tigrinus*; Schreber, 1775) (Cañon-Franco *et al.*, 2013).

There are also a few records of the presence of antibodies of this parasite in other wild neotropical carnivores. A research on neotropical mammals, included in its analysis the coati mundi (*Nasua nasua* Linnaeus, 1766) and common opossum (*Didelphis marsupialis* Linnaeus, 1758) (Thoisy *et al.*, 2003). Other carnivores such as canids have been used as indicators for *T. gondii*, however these studies have been done in the neartic regions (Dubey *et al.*, 1999). In Mexico, studies about *T. gondii* in wildlife are in small numbers and also focused on neartic species (Kikuchi *et al.*, 2004; Suzan & Ceballos, 2005; Rendón-Franco *et al.*, 2012). In areas where domestic cats are not present, wild felids are the final hosts; therefore it is very important to evaluate which role they play in the wild cycle of the parasite. In the other hand, it is important to determine if other carnivores or marsupials participate in the wild cycle of *T. gondii* as an infectious host for the felids or even for humans, since some carnivores such as the coati mundi, skunk, and opossum are consumed by people in rural areas (Naranjo *et al.*, 2004;

Tlapaya & Gallina, 2010; Contreras-Moreno *et al.*, 2012).

The objective of this study is to determine the presence of *T. gondii* antibodies in wild carnivores and marsupials in a neotropical area within northeast Mexico.

## MATERIAL AND METHODS

As a part of a felid ecological project developed by Caso, captures were done with the use of box traps (Tomahawk live Trap model No. 109.5, Tomahawk Live Trap Company, Hazelhurst, Wisconsin 54531, USA; Caso, 2013). Blood was obtained from jaguarundis (*Puma yagouaroundi* É. Geoffroy, 1803), coyotes (*Canis latrans* Say, 1823), coati mundis (*Nasua narica* Linnaeus, 1766), bobcats (*Lynx rufus* Schreber, 1777), hog-nosed skunks (*Conepatus leuconotus*; Lichtenstein, 1832), and common opossums (*Didelphis* sp.) at Los Ebanos and Los Pericos (23° 27' N, 97° 48' W) cattle ranches from 1998 to 2006 (Caso, 2013). In the other hand, captures of margays (*Leopardus wiedii* Schinz, 1821), gray foxes (*Urocyon cinereoargenteus* Schreber, 1775) and coati mundis were done at El Cielo Biosphere Reserve (23°03'N, 90°13'W) in 2003 (Carvajal-Villarreal *et al.*, 2012). Both study areas are in the state of Tamaulipas, Mexico. *P. yagouaroundi* and *L. wiedii* are protected by Mexican laws under category of near threatened and risk of extinction respectively (SEMARNAT, 2010). All carnivores in this study are considered as least concern by the IUCN, except *L. wiedii*, which is considered near threatened (IUCN, 2014).

Samples were obtained after chemical immobilization, and serum was maintained frozen at -20 C° until lab analysis (Caso *et al.*, 2005; Caso, 2013). Each individual was safely handled according to the recommendations of the American Society of Mammalogists (Sikes & Gannon, 2007) and all were released at the same capture point when the effects of immobilization were not present.

Antibody detection was done through a test latex

agglutination test (Toxotest-MT, Eiken Chemical Co. LTD, Tokyo 110-8408, Japan) accordingly with lab specifications. This test has been used before in wild felids to test the presence of *T. gondii* (Ramos *et al.*, 2001; Kikuchi *et al.*, 2004; Rendón-Franco *et al.*, 2012). Titers > 1:32 were considered positives for felines while titers > 1:16 were considered positives for the other species according with manufacture recommendations. We calculated the frequencies and confidence intervals 95% for each species using epidemiological software (Epidat 3.1 ® software, Servicio de Epidemiología Dirección Xeral de Innovación e

Xestión de Saúde Pública, Santiago de Compostela, Coruña, Spain).

## RESULTS

Results obtained were the following: margay 1 of 3 (33%), jaguarundi 0 of 2 (0%), bobcat 1 of 1 (100%), coatiundi 3 of 7 (42%), coyote 1 of 4 (25%), gray fox 2 of 3 (66%), hog-nosed skunk 0 of 1 (0%) and common opossum 1 of 5 (20%). Titers in positive animals were from 1:16 in carnivores to 1:256 in coatiundis, and positive for opossum 1:16 titer (Table 1).

Table 1. Frequencies and antibodies titer against *Toxoplasma gondii* in wild carnivores and marsupials.

Specie/serum dilution	>1:16	1:16b	1:32a	1:64	1:128	1:256	n	Positive	Frequency % (CI 95%)
Margay ( <i>Leopardus wiedii</i> )	1	1		1			3	1	33 (1-90)
Jaguarundi ( <i>Puma yagouaroundi</i> )	1	1					2	0	0 (ND)
Bobcat ( <i>Lynx rufus</i> )				1			1	1	100 (ND)
Coatiundi ( <i>Nasua narica</i> )	4	1				2	7	3	42 (10-81)
Coyote ( <i>Canis latrans</i> )	3	1					4	1	25 (1-80)
Gray fox ( <i>Urocyon cinereoargenteus</i> )	1			1	1		3	2	66 (9-99)
Hog-nosed skunk ( <i>Conepatus leuconotus</i> )	1						1	0	0 (ND)
Opossum ( <i>Didelphis</i> sp)	4	1					5	1	20 (0-72)

a=positive feline b= positive other than feline, ND=no done.

## DISCUSSION

There are not previous reports of antibodies prevalence of *T. gondii* in margay and jaguarundi; however, in jaguarundis it has been found that they can excrete oocysts naturally or during experiments (Jewell *et al.*, 1972; Pizzie *et al.*, 1978). In this study, we did not find any evidence of anti-*Toxoplasma* antibodies in jaguarundis, but because to the small sample size (n = 2), is not possible to say that jaguarundis could not be infected by the parasite in the wild. Even though the small number of individual

margays, these results are important since they ensure that margays could be infected by *T. gondii* in the wild. In the case of the gray fox, there have been reports that indicate that the prevalence of *T. gondii* in this species is high in the U.S. since it runs from 25% to 75% depending on the area, and this is consistent with the present study. Same research also found that the prevalence of *T. gondii* in coyotes is 59%, however in the present study it is lower. It is important to mention that sample size of this study for both species is low; therefore is not possible compare these results with the ones in U.S (Dubey *et al.*, 1999).

This is the study that has obtained the best sample size for wild coatimundis ( $n = 7$ ). These results indicate that the prevalence found of 42% is lower of what was found by the only previous study of 72% of prevalence (Thoisy *et al.*, 2003). Due to the small sample size in hog-nosed skunk ( $n = 1$ ), is not conclusive that they could not be infected by *T. gondii*. In the case of opossums, 20% of prevalence was record, that is within the rank (13-29%) reported for the U.S., but it is higher with the one reported (10%) for central Mexico, where mean ambient humidity is lower (Smith & Frenkel, 1995; Hill *et al.*, 1998; Suzan & Ceballos, 2005; Mitchell *et al.*, 2006). This could explain the differences since in neotropical regions there is more humidity and this helps the parasite to remain viable for longer periods. In French Guyana, it was found 15% ( $n=34$ ) of prevalence in common opossums and 20% ( $n=15$ ) in white-eared opossum (*D. albiventris*; Thoisy *et al.*, 2003). In Brazil, it was found 20.4% ( $n=396$ ) of prevalence in common opossums (Yal *et al.*, 2003).

The prevalence of *T. gondii* found in coatimundis and opossums have an important zoonotic risk, since as it was mentioned earlier; these species are hunted and consumed as food by people in rural areas of Mexico (Naranjo *et al.*, 2004; Mitchell *et al.*, 2006; Tlapaya & Gallina, 2010; Contreras-Moreno *et al.*, 2012).

The role of wild cats on the cycle of *T. gondii* is important particularly in areas where domestic cats coexist with them. In the case of other wild carnivores, there is little information about what is their role in the wild cycle of *T. gondii*. However, it is important to know their role as hosts since they compete for the infected prey and therefore they can lower the prevalence of the disease. Since opossums are commonly depredated by different carnivores including felids, it is important to determine the prevalence in them to know which is the parasite's infectious cycle. It is also important to generate studies that include intermediate hosts with the objective of identifying how the parasite cycle closes in the wild (ferraroni *et al.*, 1980).

## ACKNOWLEDGEMENTS

Thank to the Dallas Zoo, the Oklahoma City Zoo, and Los Ebanos Ranch, for their funding and support for this project. Also thank to the Dirección General de Vida Silvestre and the State of Tamaulipas to provide the necessary permits to develop this study.

## BIBLIOGRAPHIC REFERENCES

Cañon-Franco, WA, Araújo, FAP & Gennari, SM. 2013. *Toxoplasma gondii* in small neotropical wild felids. Brazilian Journal of Veterinary Research and Animal Science, vol. 50, pp. 50-67.

Carvajal-Villarreal, S, Caso, A, Downey, P, Moreno, A, Tewes, ME & Grassman Jr, LI. 2012. Spatial patterns of the margay (*Leopardus wiedii*, Felidae, Carnivora) at "El Cielo" Biosphere Reserve, Tamaulipas, Mexico. Mammalia, vol. 76, pp. 237-244.

Caso, A. 2013. Spatial differences and local avoidance of Ocelot (*Leopardus pardalis*) and Jaguarundi (*Puma yagouaroundi*) in Northeast Mexico. Tesis de Doctor en Ciencias Biológicas, Texas A&M University-Kingsville, Kingsville, Texas.

Caso, A, Carvajal-Villarreal, S, Downey, P, & Moreno, A. 2005. Técnica de captura y manejo del margay (*Leopardus wiedii*) en la Reserva de la Biosfera El Cielo. In: Sánchez-Ramos, G, Reyes-Castillo, P & Dirzo, R, (eds.). *Historia Natural de la Reserva de la Biosfera "El Cielo", Tamaulipas. México.* Universidad Autónoma de Tamaulipas, Instituto de Ecología A. C., UNAM. Toppan printing inc. Vol. 55, pp. 538-542.

Contreras-Moreno, FM, De la Cruz-Félix, K, & Bello-Gutiérrez, J. 2012. Uso patrones de cacería y preferencia de presas en dos sitios del parque estatal La Sierra, Tabasco, México. Etnobiología, vol. 10, pp. 1-9.

Della, MG. 1999. *Toxoplasmosis in zoo animals.* In: Fowler, ME, & Miller RE (eds.). *Zoo and Wild Animal Medicine*, 4<sup>th</sup> ed. W.B.

Sauders Co., Philadelphia, Pennsylvania. pp. 131-135.

Dreesen, DW. 1990. *Toxoplasma gondii infections in wildlife*. Journal of the American Veterinary Medical Association, vol. 196 pp. 274-276.

Dubey, JP, Storandt, ST, Kwok, OC, Thulliez, P & Kazacos, KR. 1999. *Toxoplasma gondii antibodies in naturally exposed wild coyotes, red foxes, and gray foxes and serologic diagnosis of toxoplasmosis in red foxes fed T. gondii oocysts and tissue cysts*. The Journal of Parasitology, vol. 85, pp. 240-243.

Ferraroni, JJ, Reed, SG & Speer, CA. 1980. *Prevalence of Toxoplasma antibodies in humans and various animals in the Amazon*. Proceedings of the Helminthological Society of Washington, vol. 47, pp. 148-150.

Frenkel, JK. 1990. *Transmission of toxoplasmosis and the role of immunity in limiting transmission and illness*. Journal of the American Veterinary Medical Association, vol. 196, pp. 233-240.

Hill, RE, Zimmerman, JJ, Wills, RW, Patton, S & Clark, WR. 1998. *Seroprevalence of antibodies against Toxoplasma gondii in free-ranging mammals in Iowa*. Journal of Wildlife Diseases, vol. 34, pp. 811-815.

Jewell, ML, Frenkel, JK, Johnson, KM, Reed, V & Ruiz, A. 1972. *Development of Toxoplasma oocysts in neotropical felidae*. The American Journal of Tropical Medicine and Hygiene, vol. 21, pp. 512-517.

Kikuchi, Y, Chomel, BB, Kasten, RW, Martenson, JS, Swift, PK & O'Brien, SJ. 2004. *Seroprevalence of Toxoplasma gondii in American free-ranging or captive pumas (Felis concolor) and bobcats (Lynx rufus)*. Veterinary Parasitology, vol. 120, pp. 1-9.

Mitchell, SM, Richardson, DJ & Lindsay, DS. 2006. *Prevalence of agglutinating antibodies to Toxoplasma gondii in striped skunks (Mephitis mephitis), opossums (Didelphis virginiana), and raccoons (Procyon lotor) from Connecticut*. The Journal of Parasitology, vol. 92, pp. 664-665.

Naranjo, EJ, Guerra, MM, Bodmer, RE & Bolaños, JE. 2004. *Subsistence hunting by three ethnic groups of the Lacandon forest, Mexico*. Journal of Ethnobiology, Vol. 24, pp. 233-253.

Pizzie, HL, Rico, CM, & Pessat, OAN. 1978. *Hallazgo del ciclo ontogenico selvatico del Toxoplasma gondii en felidos salvajes (Oncifelis geoffroyi, Felis cocolo y Felis eira) de la provincia de Cordoba*. Revista Militar de Veterinaria, vol. 25, pp. 293-300.

Ramos, SJ, Ogassawara, S, Adania, CH, Ferreira, F, Gennari, SM, & Ferreira-Neto, JS. 2001. *Seroprevalence of Toxoplasma gondii in captive neotropical felids from Brazil*. Veterinary Parasitology, vol. 102, pp. 217-224.

Rendón-Franco, E, Caso-Aguilar, A, Jiménez-Sánchez, NG, Sandoval-Sánchez, AL, Brousset, HJDM, & Zepeda-López, HM. 2012. *Prevalence of anti-Toxoplasma gondii antibody in free-ranging ocelots (Leopardus pardalis) from Tamaulipas, Mexico*. Journal of Wildlife Diseases, vol. 48, pp. 829-831.

SEMARNAT (Secretaría de Medio Ambiente y Recursos Naturales). 2010. *Norma Oficial Mexicana NOM-059-SEMARNAT-2010, protección ambiental, especies nativas de México de flora y fauna silvestre, categorías de riesgo y especificaciones para su inclusión, exclusión o cambio, lista de especies en riesgo*. Diario Oficial de la Nación. pp. 1-78.

Sikes, RS & Gannon, WL. 2007. *The animal care and use committee of the American Society of Mammalogists*. Guidelines of the American Society Mammalogists for the Use of Wild Mammals in Research. Journal of Mammalogy, vol. 88, pp. 809-23.

Smith, DD & Frenkel, JK. 1995. *Prevalence of antibodies to Toxoplasma gondii in wild mammals of Missouri and east central Kansas: biologic and ecologic considerations of transmission*. Journal of Wildlife Diseases, vol. 31, pp. 15-21.

Suzan, G & Ceballos, G. 2005. *The role of feral*

*mammals on wildlife infectious disease prevalence in two nature reserves within Mexico City limits.* Journal of Zoo and Wildlife Medicine, vol. 36, pp. 479-484.

Thoisy, B, Demar, M, Aznar, C & Carme, B. 2003. *Ecologic correlates of Toxoplasma gondii exposure in free-ranging neotropical mammals.* Journal of Wildlife Diseases, vol. 39, pp. 456-459.

Tlapaya, L & Gallina, S. 2010. *Cacería de mamíferos medianos en cafetales del centro de Veracruz, México.* Acta Zoologica Mexicana (n.s.), vol. 26, pp. 259-277.

UICN (Unión Internacional para la Conservación de la Naturaleza y los Recursos Naturales). 2014. *Red list of threatened animals.* UICN Species Survival Commission, Glad, Suiza.

Available online: <http://www.iucnredlist.org>

Wolfe, BA. 2003. *Toxoplasmosis.* In: Fowler, ME & Miller, RE (eds.). *Zoo and Wild Animal Medicine.* 5<sup>th</sup> ed. W.B. Saunders Co., St. Louis Missouri. pp. 745-749.

Yal, LEO, Cañon-Franco, WA, Geraldí, VC, Summa, MEL, Camargo, MCGO, Dubey, JP & Gennari, SM. 2003. *Seroprevalence of Neospora caninum and Toxoplasma gondii antibodies in the South American opossum (Didelphis marsupialis) from the city of São Paulo, Brazil.* The Journal of Parasitology, vol. 89, pp. 870-871.

Received August 1, 2014.  
Accepted October 21, 2014.